

# From elusive thio- and selenosilanoic acids to copper(i) complexes with intermolecular Si=E → Cu–O–Si coordination modes (E = S, Se)<sup>†</sup>

Gengwen Tan, Yun Xiong, Shigeyoshi Inoue, Stephan Enthaler, Burgert Blom, Jan D. Epping and Matthias Driess\*

Cite this: *Chem. Commun.*, 2013, 49, 5595

Received 16th March 2013,  
Accepted 29th April 2013

DOI: 10.1039/c3cc41965g

www.rsc.org/chemcomm

The facile synthesis of the first stable selenosilanoic acid–base adduct  $\text{LSi}(=\text{Se})\text{OH}(\text{dmap})$  **3** ( $\text{L} = \text{CH}[\text{C}(\text{Me})\text{NAr}]_2$ ,  $\text{Ar} = 2,6\text{-iPr}_2\text{C}_6\text{H}_3$ ,  $\text{dmap} = 4\text{-dimethylaminopyridine}$ ), the heavier analogue of the thiosilanoic acid adduct  $\text{LSi}(=\text{S})\text{OH}(\text{dmap})$  **1**, is reported. Both adducts **1** and **3** react readily with  $\text{MesCu}$  ( $\text{Mes} = 2,4,6\text{-trimethylphenyl}$ ) to form the novel dimeric  $\text{Cu}(\text{i})$  complexes  $[\text{LSi}(=\text{E})\text{OCu}]_2$  (**4**:  $\text{E} = \text{S}$ ; **5**:  $\text{E} = \text{Se}$ ) with unprecedented intermolecular  $\text{Si}=\text{E} \rightarrow \text{Cu}-\text{O}-\text{Si}$  coordination modes. The latter are efficient pre-catalysts for the  $\text{Cu}(\text{i})$ -mediated aziridination of styrene with  $\text{PhI}=\text{N}(\text{Ts})$  ( $\text{Ts} = \text{tosyl}$ ).

Using organic molecular defined species containing  $\text{M}-\text{O}-\text{Si}$  moieties as structural models for silica-supported metal/metal oxide catalysts has been proved to be a facile method to unravel the structure of a pre-catalyst as well as the mechanism of a catalytic reaction.<sup>1</sup> For example, the globular 56-membered copper(i) siloxane containing core made up of  $\text{Cu}-\text{O}-\text{Si}$  moieties (Chart 1), which is even soluble in organic solvents, could be prepared by the reaction of silanetriol with  $(\text{MesCu})_4$  ( $\text{Mes} = 2,4,6\text{-Me}_3\text{C}_6\text{H}_2$ ). This copper siloxane cluster is active for the Ullmann–Goldberg-type  $\text{C}-\text{N}$  coupling reaction,<sup>2</sup> and can serve as a structural and functional model for silica-supported copper (pre)catalysts. In order to better understand the structure–reactivity relationships of  $\text{Cu}-\text{O}-\text{Si}$  systems for various  $\text{Cu}$ -based chemical transformations, the synthesis of other types of  $\text{Cu}-\text{O}-\text{Si}$  containing compounds as structural and functional models for metallated terminal OH groups of silica surfaces is desired.

Organic silanols have been reported to form well defined  $\text{M}-\text{O}-\text{Si}$  type compounds including silanediols, disiloxane-1,3-diols, silanetriols, trisiloxane-diol and silsesquioxane-triols (Chart 1).<sup>1a,b,3,4</sup> As possible intermediates involved in silica synthesis through hydrolysis of suitable silicon(IV) starting materials under acidic conditions, silicic acids,  $\text{SiO}_x(\text{OH})_{4-2x}$ , have not received much attention as precursors for the selective synthesis of  $\text{M}-\text{O}-\text{Si}$  compounds owing to their elusive nature. Likewise, organic silanoic acids  $\text{RSi}(=\text{O})\text{OH}$ ,





Chart 1

the silicon analogues of carboxylic acids, represent elusive species which could only be studied at liquid nitrogen temperatures because of the presence of the highly polarized  $\text{Si}=\text{O}$  subunit which can undergo facile isomerisation or intermolecular head-to-tail polymerisation.<sup>5</sup> Recently, we reported the synthesis of the first isolable silanoic acid–base adducts, including the thiosilanoic system  $\text{LSi}(=\text{S})\text{OH}(\text{dmap})$  **1** ( $\text{L} = \text{CH}[\text{C}(\text{Me})\text{NAr}]_2$ ,  $\text{Ar} = 2,6\text{-iPr}_2\text{C}_6\text{H}_3$ ,  $\text{dmap} = 4\text{-dimethylaminopyridine}$ ) which is stabilised *via* hydrogen bonding to the pyridine N atom of  $\text{dmap}$  (Chart 1).<sup>6</sup> With these compounds in hand it became possible to employ **1** as a building block for  $\text{Si}(=\text{S})-\text{O}-\text{M}$  formation. Accordingly, we reported the formation and reactivity of an isolable monomeric  $\text{Si}(=\text{S})-\text{O}-\text{Mn}(\text{ii})$  complex.<sup>7</sup> Herein, we report the unexpectedly facile synthesis of the first isolable selenosilanoic acid–base adduct  $\text{LSi}(=\text{Se})\text{OH}(\text{dmap})$  **3** (Chart 1), and the  $\text{Cu}$ -metallation reactions of **1** and **3** with  $(\text{MesCu})_4$  to give the novel dimeric  $\text{LSi}(=\text{E})\text{OCu}$  complexes **4** ( $\text{E} = \text{S}$ ) and **5** ( $\text{E} = \text{Se}$ ) with unprecedented intermolecular  $\text{Si}=\text{E} \rightarrow \text{Cu}-\text{O}-\text{Si}$  coordination modes. In addition, their ability to serve as efficient pre-catalysts

Technische Universität Berlin, Department of Chemistry, Metalorganics and Inorganic Materials, Sekr. C2, Strasse des 17. Juni 135, 10623 Berlin, Germany.  
E-mail: Matthias.driess@tu-berlin.de; Fax: +49 30-314-29732

<sup>†</sup> Electronic supplementary information (ESI) available: Experimental details and crystal data and structure refinement for **3–5**. CCDC 921657–921659. For crystallographic data in CIF or other electronic format see DOI: 10.1039/c3cc41965g



**Scheme 1** Synthesis of compound 3.


in the aziridation of a C=C bond with  $\text{PhI}=\text{N}(\text{Ts})$  has been demonstrated.

The thiosilanoic acid–base complex **1** is accessible from the reaction of the corresponding stable silanone complex  $\text{L}'\text{Si}(=\text{O})(\text{dmap})$  ( $\text{L}' = \text{CH}[\text{C}(\text{Me})(\text{C}=\text{CH}_2)](\text{NAr}_2)$ ,  $\text{Ar} = 2,6\text{-iPr}_2\text{C}_6\text{H}_3$ ) with  $\text{H}_2\text{S}$  at ambient temperature. Considering the toxicity of  $\text{H}_2\text{Se}$ , we attempted to synthesize the selenium congener of **1** with dilithium selenide as a selenium source, which can be obtained from the reaction of elemental selenium with lithium triethylhydridoborate in THF.<sup>8</sup> Treatment of the *in situ* prepared dilithium selenide with one equivalent of **2** in THF at  $-78^\circ\text{C}$ , followed by protonation with two molar equivalents of trimethylammonium chloride, leads to a clear yellow solution, from which compound **3** could be isolated in the form of yellow crystals in 75% yield (Scheme 1).


The strikingly stable compound **3** has very similar solubility properties to **1**. It is soluble in toluene, THF, chloroform, dichloromethane, marginally soluble in benzene but insoluble in *n*-hexane. It was fully characterised using multinuclear NMR spectroscopy, elemental analysis, mass spectrometry and IR spectroscopy, as well as single-crystal X-ray diffraction analysis. The high resolution electrospray ionization mass spectrometry (ESI-MS) shows the molecular ion peak at  $m/z$  665.3138 [ $\text{LSi}(=\text{Se})\text{OH}(\text{DMAP}) + \text{H}$ ]<sup>+</sup> as the base signal. In the  $^1\text{H}$  NMR spectrum in  $\text{CDCl}_3$ , the chemical shift of the ring proton in the  $\gamma$ -position of the  $\beta$ -diketiminato L is observed at  $\delta = 5.73$  ppm. This chemical shift is comparable to that observed for **1** ( $\delta = 5.71$  ppm). The proton of the OH group resonates at  $\delta = 6.42$  ppm in the same region as the proton resonance signals for the pyridine moiety of dmap. The  $^{29}\text{Si}\{^1\text{H}\}$  resonance signal of **3** in  $\text{CDCl}_3$  at  $\delta = -25.5$  ppm is low-field shifted compared to that of **1** ( $\delta = -30.0$  ppm), while its  $^{77}\text{Se}\{^1\text{H}\}$  NMR spectrum exhibits a singlet at  $\delta = -545.2$  ppm, which is up-field shifted compared to those of the related selenosilanoic silyl ester diastereomers ( $\delta = -384.8$  and  $-401.3$  ppm).<sup>9</sup>

The molecular structure of compound **3** is depicted in Fig. 1. It crystallizes in the triclinic space group  $P\bar{1}$  with one *n*-hexane molecule lying about an inversion center. It is isotopic with **1**, with one dmap ligand connected to the selenosilanoic acid moiety through an O–H $\cdots$ N hydrogen bond. The Si1–O1 bond length of 1.619(2) Å is comparable to that of **1** (1.620(2) Å), whereas the Si(1)–Se(1) bond distance of 2.1348(7) Å is close to the Si=Se distance of a related selenosilanoic silyl ester (2.117(1) Å).<sup>9</sup> As expected, other metric parameters of **3** are akin to those of **1**.

Both compounds **1** and **3** were allowed to react with  $(\text{MesCu})_4$  ( $\text{Mes} = 2,4,6\text{-trimethylphenyl}$ ) in the hope of producing the corresponding Cu(I) complexes **4** and **5**, respectively.  $(\text{MesCu})_4$  is




**Fig. 1** Molecular structure of compound **3** in the solid state with the 50% probability level for the core structure. Hydrogen atoms (except for H1) and the *n*-hexane molecules are omitted for the sake of clarity.




**Scheme 2** Syntheses of the Cu(I) complexes **4** and **5**.

known to serve as a smooth reagent for preparing Cu(I) complexes by deprotonation of amines and silanols.<sup>2,3,10</sup>

Accordingly, the reaction of compound **1** with  $\frac{1}{4}$  molar equivalent of  $(\text{MesCu})_4$  was carried out in THF at  $-20^\circ\text{C}$ . The  $^1\text{H}$  NMR spectrum of the resulting reaction mixture already shows the absence of the resonance signals of the OH group of **1** and those of  $(\text{MesCu})_4$  with concomitant liberation of dmap, indicating that the metallation of **1** was successful. Indeed, the desired compound **4** could be isolated from the reaction mixture as yellow crystals in 85% yield (Scheme 2). Single crystals of **4** suitable for X-ray diffraction analysis could be obtained in toluene solutions at  $0^\circ\text{C}$ . The structure analysis revealed that the compound is a dimer with intermolecular Si=S  $\rightarrow$  Cu–O–Si interactions (Fig. 2). The compound possesses  $C_2$  symmetry and consists of a planar eight-membered  $\text{Si}_2\text{O}_2\text{S}_2\text{Cu}_2$  ring with two *trans*-oriented  $\beta$ -diketiminato ligands L. The Cu centres are linearly coordinated by one oxygen and a sulfur atom of the neighbouring Si=S subunit. The Cu(1) $\cdots$ Cu(1A) distance of 2.8135(7) Å suggests a weak  $d^{10}\cdots d^{10}$  interaction in this molecule.<sup>11</sup> The Si(1)–O(1) distance of 1.568(2) Å is significantly shorter than that in the precursor **1** (1.620(2) Å), whereas the Si(1)–S(1) bond length (2.0609(9)) is longer than that



**Fig. 2** (a) Molecular structures of compounds **4** (E=S) and **5** (E=Se) in the solid state with the 50% probability level for the core structure. Hydrogen atoms and the toluene molecules are omitted for the sake of clarity. Operation symmetry for all atoms labelled "A":  $-x + 3/2, -y + 1/2, -z$  (**4**);  $-x + 2, -y, -z$  (**5**); (b) representation of the eight-membered planar  $\text{Si}_2\text{O}_2\text{S}_2\text{Cu}_2$  ring without substituents at silicon atoms.



**Fig. 3** Optimized structure for the proposed stereoisomer **4b**. Hydrogen atoms are omitted for the sake of clarity.

observed in **1** (1.993(1) Å), suggesting carboxylate-like  $\pi$ -conjugation in the SiSO moiety.<sup>12</sup>

The composition of **4** has been confirmed by multinuclear NMR spectroscopy, elemental analysis and IR spectroscopy. The solubility of **4** is similar to that of precursor **1**. Unexpectedly, two sets of resonances for the  $\beta$ -diketiminato ligand **L** appear in the <sup>1</sup>H NMR spectrum with a ratio of 1 : 0.62 as indicated by the integrals of the resonances from the ring proton in the  $\gamma$ -position of **L** at  $\delta$  = 5.55 and 5.49 ppm, respectively. Accordingly, the <sup>29</sup>Si{<sup>1</sup>H} NMR spectrum reveals two close signals at  $\delta$  = -38.0 and -39.1 ppm, respectively. The two sets of resonances suggest the presence of two stereoisomers of **4** (**4a** and **4b**) in chloroform solutions. Indeed this is substantiated by the results obtained from diffusion ordered spectroscopy (DOSY) experiments, revealing identical diffusion coefficients for these two species (see ESI<sup>†</sup>) and thus the same molecular size and composition, respectively. In other words, dissociation of **4** in chloroform solutions can be excluded. As expected, the two stereoisomers can be interconverted as shown using variable <sup>1</sup>H NMR spectroscopy. Cooling of a CDCl<sub>3</sub> solution of **4** to 230 K changes the ratio of signal sets of **4a** and **4b** from 1 : 0.62 at ambient temperature to 1 : 0.42 (see ESI<sup>†</sup>). In the solid state <sup>29</sup>Si NMR spectrum of **4** in crystalline form, there is only one resonance signal at  $\delta$  = -41.6 ppm, whereas in the spectrum of the fine powder two signals are observed ( $\delta$  = -41.6 ppm and -38.7 ppm) (see ESI<sup>†</sup>), this suggests that **4a** holds a structure as shown in Fig. 2 with two ligands **L** in the *trans*-position. We inferred that the other stereoisomer **4b** preserves the *C*<sub>2</sub> symmetry of **4a** but the two ligands **L** are now *cis*-oriented. However, this is ruled out by the results of DFT calculations, which revealed that the proposed *cis* isomer is least favoured. Instead the DFT calculations suggest a twisted Si<sub>2</sub>O<sub>2</sub>Se<sub>2</sub>Cu<sub>2</sub> core structure also with *C*<sub>2</sub> symmetry as a stereoisomer which is only 5.5 kJ mol<sup>-1</sup> less stable than **4a** (Fig. 3 and ESI<sup>†</sup>).

Similar to the synthesis of **4**, compound **5** could be obtained by the reaction of **3** with (MesCu)<sub>4</sub> in THF (Scheme 2). Compound **5** has been fully characterized by multinuclear NMR spectroscopy, elemental analysis, IR and mass spectroscopy, and X-ray crystallography. Akin to the situation of **4**, there are two stereoisomers present in CDCl<sub>3</sub> solutions as shown by the <sup>1</sup>H, <sup>13</sup>C, <sup>29</sup>Si NMR spectra as well as <sup>1</sup>H-DOSY experiments (see ESI<sup>†</sup>).

Single-crystals of **5** in the triclinic space group *P* $\bar{1}$  could be obtained in toluene solutions. The X-ray diffraction analysis revealed that **5** and **4** are isotopic (Fig. 2). Akin to the structure of **4**, compound **5** is a dinuclear copper(i) complex with both copper centers coordinated by O and Se atoms. A linear geometry of the O–Cu–Se connection with an angle of 172.07(7) $^\circ$  is observed. The Cu(1)···Cu(1A) bond distance of 2.9271(8) Å is slightly longer than that in compound **4** (2.8135(7) Å). This can be explained by

the longer Si–Se distance (2.2011(9) Å) compared to that of the Si–S bond (2.0609(9) Å) in **4**.

It has been shown that Cu(i) complexes can be efficiently applied as pre-catalysts in metal-catalyzed nitrene-transfer reactions if the Cu(i) centre is efficiently chelate coordinated.<sup>13</sup> In a preliminary study the catalytic ability of compounds **4** and **5** in the nitrene-transfer reaction (aziridation) of a C=C bond has been evaluated. The catalytic reactions were carried out by using styrene and PhI=N(Ts) (Ts = tosyl) as a nitrene source in the presence of 2.5 mol% of **4** and **5**, respectively, in CH<sub>2</sub>Cl<sub>2</sub> at ambient temperature. The resulting yields of the *N*-tosyl-2-phenylaziridine product [85% (**4**) and 87% (**5**)] are similar to a reported result.<sup>13a</sup> It is generally considered that a Cu(i)–nitrene species is the active component for the aziridation reaction.<sup>13b</sup> Thus, it is reasonable to assume that the dimeric Cu(i) complexes react initially with PhI=N(Ts) to form the corresponding Cu(i)–nitrene intermediates, which are capable of facile nitrene-transfer to the C=C bond of styrene.

In summary, using dilithium selenide as a selenium source, the first isolable selenosilanoic acid–base adduct **3** has been synthesized which is isostructural with the thiosilanoic acid–base adduct **1**. Facile reaction of **1** and **3** with  $\frac{1}{4}$  molar equivalents of (MesCu)<sub>4</sub> led to the unprecedented dimeric copper(i) complexes **4** and **5**. Both complexes exist in two stereoisomeric forms in chloroform solutions and verify the novel Si=E  $\rightarrow$  Cu–O–Si structural motif in Cu(i) siloxane chemistry. In addition, compounds **4** and **5** can act as reliable pre-catalysts in aziridation of styrene with PhI=N(Ts).

## Notes and references

- (a) R. Murugavel, A. Voigt, M. G. Walawalkar and H. W. Roesky, *Chem. Rev.*, 1996, **96**, 2205–2236; (b) R. Murugavel, V. Chandrasekhar and H. W. Roesky, *Acc. Chem. Res.*, 1996, **29**, 183–189; (c) N. Winkhofer, A. Voigt, H. Dorn, H. W. Roesky, A. Steiner, D. Stalke and A. Reller, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 1352–1354; (d) H. C. L. Abbenhuis, *Chem.–Eur. J.*, 2000, **6**, 25–32; (e) K. L. Furdala, I. J. Drake, A. T. Bell and T. D. Tilley, *J. Am. Chem. Soc.*, 2004, **126**, 10864–10866.
- G. Tan, Y. Yang, C. Chu, H. Zhu and H. W. Roesky, *J. Am. Chem. Soc.*, 2010, **132**, 12231–12233.
- F. Schax, C. Limberg and C. Mügge, *Eur. J. Inorg. Chem.*, 2012, 4661–4668.
- (a) V. Lorenz, A. Fischer, K. Jacob, W. Brueser and F. T. Edelmann, *Chem.–Eur. J.*, 2001, **7**, 848–857; (b) V. Lorenz, S. Gießmann, Y. K. Gun'ko, A. K. Fischer, J. W. Gilje and F. T. Edelmann, *Angew. Chem., Int. Ed.*, 2004, **43**, 4603–4606; (c) V. Lorenz, A. Fischer, K. Jacob, W. Brueser, T. Gelbrich, P. G. Jones and F. T. Edelmann, *Chem. Commun.*, 1998, 2217–2218.
- (a) C. A. Arrington, R. West and J. Michl, *J. Am. Chem. Soc.*, 1983, **105**, 6176–6177; (b) R. Withnall and L. Andrews, *J. Am. Chem. Soc.*, 1985, **107**, 2567–2568.
- Y. Xiong, S. Yao and M. Driess, *Angew. Chem., Int. Ed.*, 2010, **49**, 6642–6645.
- S. Yao, Y. Xiong and M. Driess, *Chem.–Eur. J.*, 2012, **18**, 11356–11361.
- U. Herzog, H. Lange, H. Borrmann, B. Walfort and H. Lang, *J. Organomet. Chem.*, 2004, **689**, 4909–4916.
- S. Yao, Y. Xiong, M. Brym and M. Driess, *Chem.–Asian. J.*, 2008, **3**, 113–118.
- (a) G. Tan and H. Zhu, *Inorg. Chem.*, 2011, **50**, 6979–6986; (b) M. Stollenz and F. Meyer, *Organometallics*, 2012, **31**, 7708–7727.
- (a) P. K. Mehrotra and R. Hoffmann, *Inorg. Chem.*, 1978, **17**, 2187–2189; (b) K. M. Merz Jr. and R. Hoffmann, *Inorg. Chem.*, 1988, **27**, 2120–2127.
- (a) R. D. Mounts, T. Ogura and Q. Fernando, *Inorg. Chem.*, 1974, **13**, 802–805; (b) A. S. Filatov, O. Hietsoi, Y. Sevryugina, N. N. Gerasimchuk and M. A. Petrukhina, *Inorg. Chem.*, 2010, **49**, 1626–1633.
- (a) P. J. Pérez, M. Brookhart and J. L. Templeton, *Organometallics*, 1993, **12**, 261–262; (b) Y. M. Badiei, A. Krishnaswamy, M. M. Melzer and T. H. Warren, *J. Am. Chem. Soc.*, 2006, **128**, 15056–15057.