

Binuclear Manganese(III) Complexes as Electron Donors in D1/D2/Cytochrome *b*559 Preparations Isolated from Spinach Photosystem II Membrane Fragments

S. I. Allakhverdiev^{a,b,c}, M. S. Karacan^b, G. Somer^b, N. Karacan^b,
E. M. Khan^c, S. Y. Rane^c, S. Padhye^c, V. V. Klimov^a and G. Renger^d

^a Institute of Soil Science and Photosynthesis, RAS, Pushchino, Moscow Region, 142292 Russia

^b Department of Chemistry, University of Gazi, Ankara, 06500, Turkey

^c Department of Chemistry, University of Poona, 411007, India

^d Institut für Biophysikalische und Physikalische Chemie, Technische Universität, D-10623 Berlin, Bundesrepublik Deutschland

Dedicated to Professor Achim Trebst on the occasion of his 65th birthday

Z. Naturforsch. **49c**, 587–592 (1994); received June 24/July 19, 1994

PS II Reaction Center, Manganese, Pheophytin Photoaccumulation, NADP⁺ Reduction

The capability of different manganese complexes to act as PS II electron donors in D1/D2/cytochrome *b*559 complexes has been analyzed by measuring actinic light-induced absorption changes at 680 nm (650 nm) and 340 nm, reflecting the photoaccumulation of Pheophytin[−] (Pheo[−]) and the reduction of NADP⁺, respectively. The data obtained reveal: a) the donor capacity of synthetic binuclear Mn(III)₂ complexes containing aromatic ligands significantly exceeds that for MnCl₂ in both cases, *i.e.* Pheo[−] photoaccumulation and NADP⁺ reduction; b) manganese complexes can serve as suitable electron donors for light-induced NADP⁺ reduction catalyzed by D1/D2/cytochrome *b*559 complexes and ferredoxin plus ferredoxin-NADP⁺ reductase under anaerobic conditions and c) the specific turnover rate of the system leading to NADP⁺ reduction is extremely small.

The implications of these findings are briefly discussed.

Introduction

The essential steps of photosynthetic water cleavage take place within an integral membrane complex associated with extrinsic regulatory polypeptides. This operational unit is referred to as photosystem II (PS II) complex. The primary reactions of PS II comprise light absorption, excitation energy transfer to the photoactive pigment (a special chlorophyll *a* designated as P680) and electron transfer from its excited singlet state to pheophytin *a* (Pheo) acting as primary acceptor and subsequent stabilization of the primary charge separation by rapid electron transfer from Pheo[−] to a special plastoquinone molecule Q_A (Renger, 1992). Based on similarities of the functional and structural organization of the primary processes, components P680, Pheo and Q_A were inferred to

be incorporated into a heterodimer of polypeptides D1 and D2 of PS II in an analogous way as the corresponding groups are incorporated into the heterodimer of the L- and M-subunit of the reaction centers of purple bacteria (Michel and Deisenhofer, 1988; Trebst, 1986). This idea is strongly supported by the isolation of PS II complexes that i) contain only D1 and D2 together with cytochrome *b*559 (= Cyt *b*559, two subunits) and at least one smaller polypeptide, and ii) retain the ability to perform the primary charge separation (Nanba and Satoh, 1987; Barber *et al.*, 1987). Although deprived of Q_A and the oxygen-evolving complex (OEC), the D1/D2/cytochrome *b*559 complexes were found to catalyze the electron transport from an artificial electron donor (diphenylcarbazide) to an exogenous electron acceptor like silicomolybdate (Chapman *et al.*, 1988; Takahashi *et al.*, 1989). This observation implies that an electron transfer can take place from Pheo[−] to exogenous acceptors and that either this reaction or the electron donation are able competing with the rapid internal charge recombina-

Reprint requests to Prof. Dr. Gernot Renger, Technische Universität Berlin, Max-Volmer-Institut für Biophysikalische und Physikalische Chemie, Straße des 17. Juni 135, D-10623 Berlin, Bundesrepublik Deutschland.
Telefax: 49-30-31 42 11 22.

tion between Pheo⁻ and P680⁺. As the redox potential of Pheo/Pheo⁻ is rather low ($E_{m,7} = -610$ mV, see Klimov *et al.*, 1979), it appeared worthwhile to check if D1/D2/cytochrome *b*559 preparations are also able to reduce NADP⁺ ($E_{m,7} = -320$ mV). Recent reports claim that this can be the case (Allakhverdiev and Klimov, 1990, 1992; Arnon and Barber, 1990; Klimov *et al.*, 1986). In the present study manganese complexes were tested as electron donors for photoaccumulation of Pheo⁻ in the absence of exogenous electron acceptors and for NADP⁺ reduction under anaerobic conditions in the presence of ferredoxin (Fd) and Fd-NADP⁺ reductase. The results obtained indicate that synthetic binuclear complexes are efficient electron donors in D1/D2/cytochrome *b*559 preparations.

Materials and Methods

D1/D2/cytochrome *b*559 preparations from spinach were obtained from digitonin/Triton X-100 PS II fractions (DT 20 preparations, see Allakhverdiev *et al.*, 1992; Klimov *et al.*, 1982) according to the method of Nanba and Satoh (1987) with some modifications as described in Allakhverdiev *et al.* (1992). The pigment content of the isolated D1/D2/cytochrome *b*559 was determined by HPLC. A ratio of about six chlorophylls per two pheophytins was found.

A single-beam differential spectrophotometer with a phosphoroscope similar to that described previously (Klimov *et al.*, 1979, 1986; Allakhverdiev and Klimov, 1990, 1992; Allakhverdiev *et al.*, 1992) was used to monitor at 650 nm and 680 nm

the light-induced photoaccumulation of pheophytin⁻ (= Pheo⁻). Likewise, at 340 nm the photo-reduction of NADP⁺ was measured. Optical pathlength: 1 cm; measuring light beam $\lambda = 340$ nm, 650 nm or 680 nm; $I \approx 0.15 \text{ W} \cdot \text{m}^{-2}$; actinic light: $\lambda > 600$ nm, $I \approx 100 \text{ W} \cdot \text{m}^{-2}$. All measurements were performed at 20 °C.

The binuclear complexes $[\text{Mn(III)Mn(III)(HNQOX)}_4(\text{OAc})_2]$ and $[\text{Mn(III)-O-Mn(III)(HNQOX)}_2(\text{OAc})_2(\text{H}_2\text{O})_2]$ symbolized by M-2 and M-3, respectively, contained 2-hydroxy-1,4-naphthoquinone monoxime (HNQOX) as terminal and CH_3COO^- (OAc) as bridging ligands (in M-3 there exist in addition a μ -oxo-bridge). These complexes were synthesized according to previously described procedures (Charles, 1963; Wiegardt *et al.*, 1985) and characterized by spectroscopic and electrochemical methods (Khan, 1993). Ferredoxin and ferredoxin-NADP⁺ reductase were purchased from Sigma.

The compositions of the sample suspensions used in the different experiments are given in the figure legends.

Results and Discussion

When thylakoid and PS II membrane fragments are exposed to actinic illumination in the presence of the strong reductant dithionite ($\text{Na}_2\text{S}_2\text{O}_4$), Pheo⁻ accumulates and gives rise to a marked quenching of the chlorophyll fluorescence (Allakhverdiev and Klimov, 1990, 1992; Allakhverdiev *et al.*, 1978, 1979, 1986; Shuvalov *et al.*, 1980; Renger and Kaye, 1987). Analogously, Pheo⁻ photoaccumulation can be achieved also in




Fig. 1. Actinic light-induced absorption changes at 650 nm and 680 nm as a function of time in D1/D2/cytochrome *b*559 complexes. The sample suspension contained: D1/D2/cytochrome *b*559 complexes (5 μg chlorophyll/ml), 35 mM NaCl, 5 mM MgCl_2 , 1 μM methylviologen, 20 mM tris(hydroxymethyl)aminoethane-HCl, pH = 8.0. Arrows indicate turning on (up) and off (down) the actinic light. Trace 1: control with no further addition; trace 2: addition of 1 mg $\text{Na}_2\text{S}_2\text{O}_4$ /ml; trace 3: addition of 50 μM MnCl_2 , traces 4a/b: addition of 5 μM manganese complex M-3.

D1/D2/cytochrome *b*559 preparations in the presence of methylviologen (MV) and $\text{Na}_2\text{S}_2\text{O}_4$ (Barber *et al.*, 1987; Nanba and Satoh, 1987; Allakhverdiev *et al.*, 1992; Shuvalov *et al.*, 1989) and monitored by characteristic absorption changes in the red region. Light-induced absorption changes of D1/D2/cytochrome *b*559 preparations are shown in Fig. 1. In the absence of $\text{Na}_2\text{S}_2\text{O}_4$ /MV virtually no absorption change can be seen at 682 nm (trace 1 of Fig. 1), a wavelength characteristic for Pheo⁻ formation (Allakhverdiev *et al.*, 1992; Klimov *et al.*, 1979). Lack of light-induced absorption changes is understandable because the primary radical pair recombines *via* kinetics in the time domain of tens of nanoseconds (Crystall *et al.*, 1989; Danelius *et al.*, 1987; Takahashi *et al.*, 1987). On the other hand, in the presence of $\text{Na}_2\text{S}_2\text{O}_4$ /MV the actinic light causes a strong bleaching at 682 nm that biphasically relaxes in the subsequent dark period (trace 2 of Fig. 1). This effect is understandable by electron donation to P680⁺ competing with the fast recombination reaction by Pheo⁻. For kinetic reasons the quantum efficiency of this process is rather low (for a numerical analysis in thylakoids, see ref. Renger and Kaye, 1987). Pheo⁻ photoaccumulation therefore provides an indirect test for the efficiency of PS II donors. Accordingly, this assay was used to analyze several manganese compounds for their capacity to act as PS II donor in D1/D2/cytochrome *b*559 preparations. Trace 3 of Fig. 1 reveals that a much less pronounced bleaching is observed if $\text{Na}_2\text{S}_2\text{O}_4$ is substituted by 50 μM MnCl_2 acting as exogenous PS II electron donor (trace 3 of Fig. 1). At this concentration, the MnCl_2 to P680 ratio is about 60 based on a Chl content of 6 molecules per two pheophytins in the D1/D2/cytochrome *b*559 preparations used in this study. In contrast to the comparatively small Pheo⁻ photoaccumulation achieved by MnCl_2 even at excess concentrations, a full restoration of this activity is achieved by addition of the binuclear manganese complex $[\text{Mn}_2\text{O}(\text{LwOx})_2(\text{OAc})_2(\text{H}_2\text{O})_2]$ where LwOx symbolizes the ligand 2-hydroxy-1,4-naphthoquinone monoxime and OAc is acetate. The amount of this complex (referred to as M-3) required to elicit the same extent of 682 nm bleaching as achieved by 1 mg/ml $\text{Na}_2\text{S}_2\text{O}_4$ (compare traces 2 and 4a in Fig. 1) is comparatively small

(10–12 manganese per P680). No further increase of the signal amplitude is observed at tenfold M-3 concentrations (data not shown). This effect clearly shows that compound M-3 acts as an efficient electron donor to D1/D2/cytochrome *b*559 preparations. In order to check that the photobleaching at 682 nm mediated by compound M-3 really reflects Pheo⁻ formation, comparative measurements were performed at 650 nm (trace 4b of Fig. 1) where the difference spectrum exhibits a positive band (Allakhverdiev *et al.*, 1992; Klimov *et al.*, 1978, 1979). The amplitude ratio $\Delta\Delta_{650}/\Delta\Delta_{682}$ of the data nicely fits with that of the difference spectrum of Pheo⁻ *versus* Pheo.

In a recent study (Allakhverdiev *et al.*, 1994) evidence was presented that the binuclear manganese complexes M-2 and M-3 are much more efficient than MnCl_2 in electron donation to PS II and restoration of the water oxidase activity in PS II preparations that are almost completely deprived of their endogenous manganese by N,N,N',N'-tetramethyldiamine (TEMED) treatment (Ananyev *et al.*, 1992). A comparison of these findings with the data presented in Fig. 1 suggests that similar structural determinants are likely to be responsible for the different efficiencies of MnCl_2 and M-3 (M-2) in PS II complexes and D1/D2/cytochrome *b*559 preparations. Accordingly, the results of Fig. 1 (traces 2 and 4a/b) can be considered as further evidence for a special affinity of D1/D2/cytochrome *b*559 complexes to manganese depending on the structure of these compounds. The possible relevance of the present data in supporting the role of amino acid residues of polypeptide D1 in the photoassembly of the tetranuclear manganese cluster in the intact water oxidase (Nixon *et al.*, 1992) remains to be clarified in future studies. Regardless of the latter problem, the data of Fig. 1 indicate the potential donor efficiency of the binuclear manganese complex M-3 in D1/D2/cytochrome *b*559 preparations.

Within the context of the donor efficiency of several compounds in D1/D2/cytochrome *b*559 complexes another phenomenon will be briefly analyzed in this study. Recently, evidence was presented for light-induced NADP^+ reduction catalyzed by D1/D2/cytochrome *b*559 preparations in the presence of ferredoxin (Fd), Fd- NADP^+ reductase, plastocyanin and the PS II donor diphenylcarbazide (Arnon and Barber, 1990). Therefore,

it appeared worth analyzing the capacity of M-3 to support NADP⁺ photoreduction in D1/D2/cytochrome *b*559 preparations. Experiments with different manganese compounds were performed under anaerobic conditions (glucose/glucose oxidase) in the presence of Fd and Fd-NADP⁺ reductase. The process was monitored at 340 nm where the NADP⁺/NADPH difference spectrum exhibits a pronounced peak. Fig. 2 shows the data obtained. In the absence of an exogenous electron donor virtually no change of the 340 nm absorption is induced by excitation of the sample with actinic light. A significant but minor increase of this absorption is observed upon illumination of D1/D2/cytochrome *b*559 preparations in the presence of 80 μ M MnCl₂ corresponding to about 50 donor molecules per P680 (trace b of Fig. 2). The slope of the absorption increase progressively declines during the exposure to actinic illumination. After cessation of the light a decrease of the 340 nm absorption change is observed. A markedly more pronounced light-induced increase of the absorbance at 340 nm is elicited by the binuclear manganese complexes M-2 and M-3 (trace c and d, respectively, in Fig. 2). In order to




Fig. 2. Actinic light-induced absorption changes at 340 nm as a function of time in D1/D2/cytochrome *b*559 complexes. The sample suspension contained: D1/D2/cytochrome *b*559 complexes (10 μ g Chl/ml), 35 mM NaCl, 3 mM MgCl₂, 3 mM NADP⁺, 10 mM ferredoxin, 6 μ g/ml Fd-NADP⁺ reductase, 10 mM glucose, 50 activity units/ml glucose oxidase, 1000 activity units/ml catalase and 20 mM Tris-HCl, pH = 8.0. Arrows indicate turning on (up) and off (down) the actinic light. Trace a: no further additions; trace b: plus 80 μ M MnCl₂; trace c: plus 10 μ M binuclear manganese complex M-2; trace d: plus 10 μ M binuclear manganese complex M-3.

confirm that these 340 nm changes are really due to NADP⁺ reduction, the light-induced difference spectra were measured in the range from 300 to 400 nm. As expected, no light-induced change is observed in the control without any addition (trace a of Fig. 3). After addition of NADP⁺ only a small increase of the absorbance is seen in the range from 300 to 400 nm (trace 2 of Fig. 3). In marked contrast, a pronounced difference spectrum arises in the presence of NADP⁺ and compound M-3. The spectral feature is characteristic for NADP⁺ reduction.

The idea of a PS II-catalyzed NADP⁺ reduction is further supported by the finding that a special dinitrobenzene derivative recently discovered to act as electron transport inhibitor of D1/D2/cytochrome *b*559 preparations (Allakhverdiev *et al.*, 1989) completely suppressed the M-3-mediated light-induced 340 nm absorption change (data not shown).

The findings reported in this study unambiguously show that M-2 and M-3 are potent mediators of light-induced NADP⁺ reduction in D1/D2/cytochrome *b*559 complexes. As in the case of photoactivation of the water oxidase activity in manganese-depleted DT20 PS II preparations the binuclear complex M-3 is also more efficient than M-2 in supporting NADP⁺ reduction in D1/D2/cytochrome *b*559 samples (compare traces c and d




Fig. 3. Extent of actinic light-induced absorption changes as a function of wavelength in D1/D2/cytochrome *b*559 complexes. Experimental conditions as in Fig. 2 but without NADP⁺. Trace a: no addition; trace b: plus 3 mM NADP⁺; trace c: plus 3 mM NADP⁺ and 10 μ M complex M-3.

of Fig. 2). Two conclusions can be drawn from the result of this study: a) the donor capacity of manganese compounds depends on their chemical nature, *i.e.* binuclear manganese complexes are much more efficient than MnCl_2 , and b) D1/D2/cytochrome *b*559 complexes can catalyze NADP^+ reduction with PS II donors under anaerobic conditions.

At first glance, the latter effect seems to be spectacular. However, it has to be emphasized that the absolute rate of NADP^+ reduction is rather slow. An evaluation of the data presented in Fig. 2 reveals that at most a rate of $2 \mu\text{mol NADPH/mg Chl} \cdot \text{h}$ can be achieved under our experimental conditions. This value has to be compared with the more than 100-fold rates obtained with silicomolybdate as electron acceptor and either 1 mM DPC (about $900 \mu\text{eq/mg Chl}$) or 1 mM MnCl_2 ($400\text{--}500 \mu\text{eq/mg Chl} \cdot \text{h}$) as PS II donors (Chapman *et al.*, 1988). Therefore, the electron transfer from the acceptor side of D1/D2/cytochrome *b*559 preparations to the enzymatic system consisting of ferredoxin and Fd- NADP^+ reductase appears to be extremely slow. Based on a NADP^+ reduction rate of $2\text{--}4 \mu\text{eq/mg Chl} \cdot \text{h}$ and a chlorophyll content of 6 per D1/D2/cytochrome *b*559 complex a rough estimation leads to values for the electron transfer time of the rate-limiting step that are of the order of 2–3 min. This phenomenon could be simply explained within the framework of the classical Marcus theory of thermally activated non-adiabatic electron transfer (Marcus and Sutin, 1985) by assuming a comparatively large distance between Pheo^- and the iron sulfur cluster of Fd. Using the empirical relation recently derived for the distance dependence of

the matrix element for electron coupling between redox groups in proteins (Moser *et al.*, 1992) distances are obtained in the range of 25–30 Å. These values are reasonable if one takes into account that the enzymes and the D1/D2/cytochrome *b*559 complexes form aggregates that are not favourable for electron transfer.

Therefore, it is concluded, that in principle Pheo^- of PS II can provide the reducing power for NADP^+ reduction. However, for kinetic reasons owing to structural constraints only marginally small rates can be achieved. It has to be emphasized that this extremely low activity was observed under optimal conditions where Pheo^- is photoaccumulated in its reduced state (see Fig. 1). In a PS II complex with open reaction center the Pheo^- becomes reoxidized by Q_A with a reaction time of about 300 ps (Bernarding *et al.*, 1994; Eckert *et al.*, 1988; Nuijs *et al.*, 1986). Therefore, the probability of an electron transfer to NADP^+ is virtually zero, *i.e.* this reaction does practically not occur under normal conditions. For theoretical reasons it might be interesting to clarify whether or not the conditions for an electron transfer from Pheo^- to NADP^+ can be drastically improved (*e.g.* by using lipophilic redox mediators).

Acknowledgements

S. I. A. is grateful to TUBITAK (Turkish Scientific and Research Council) for a grant which supported part of this study. G. R. gratefully acknowledges financial support by Deutsche Forschungsgemeinschaft (438 113/138). Part of this work was also supported by the Russian Foundation of Basic Research and by The International Science Foundation (grant MTZ 000).

- Allakhverdiev S. I., Zharmukhamedov S. K., Klimov V. V., Vasil'ev S. S., Korvatovsky B. N. and Pashchenko V. Z. (1989), Effect of dinoseb and other phenolic compounds on fluorescence decay kinetics of photosystem II chlorophyll in higher plants. *Biol. Membr.* **6**, 1147–1153.
- Allakhverdiev S. I. and Klimov V. V. (1990), Photo-reduction of NADP^+ in subchloroplast preparations of photosystem II in higher plants. *Biol. Membr.* **7**, 497–505.
- Allakhverdiev S. I. and Klimov V. V. (1992), Photo-reduction of NADP^+ in photosystem II of higher plants: Requirements for manganese. *Z. Naturforsch.* **47c**, 57–62.

- Allakhverdiev S. I., Kozlov Y. N., El-Sheekh M. M., Demeter S. and Klimov V. V. (1992), Effect of the chemical modification of tyrosine and histidine in isolated PS II reaction centers on the thermoluminescence band TL-55. *Biol. Membr.* **9**, 904–914.
- Allakhverdiev S. I., Karacan M. S., Somer G., Karacan N., Khan E. M., Rane S. Y., Padhye S., Klimov V. V. and Renger G. (1994), Reconstitution of the water-oxidizing complex in manganese-depleted photosystem II complexes by using synthetic binuclear manganese complexes. *Biochemistry* **33** (in press).
- Ananyev G., Wydrzynski T., Renger G. and Klimov V. V. (1992), Transient peroxide formation by the manganese-containing redox active donor side of

- photosystem II upon inhibition of O₂ evolution with lauroylcholine chloride. *Biochim. Biophys. Acta* **1110**, 303–311.
- Arnon D. I. and Barber J. (1990), Photoreduction of NADP⁺ by isolated reaction centers of photosystem II: Requirement for plastocyanin. *Proc. Natl. Acad. Sci. U.S.A.* **87**, 5930–5934.
- Barber J., Chapman D. J. and Telfer A. (1987), Characterization of a PS II reaction centre isolated from the chloroplasts of *Pisum sativum*. *FEBS Lett.* **220**, 67–73.
- Bernarding J., Eckert H.-J., Eichler H.-J., Napiwotzki A. and Renger G. (1994), Kinetic studies on the stabilization of the primary radical pair P680⁺Pheo⁻ in different photosystem II preparations from higher plants. *Photochem. Photobiol.* **59**, 566–573.
- Chapman D. J., Gounaris K. and Barber J. (1988), Electron-transport properties of the isolated D1/D2/cytochrome *b*559 photosystem II reaction center. *Biochim. Biophys. Acta* **933**, 423–431.
- Charles R. G. (1963), Acetylacetonatomanganese(III), in: *Inorganic Synthesis*, Vol. 7 (J. Kleinberg, ed.). McGraw Hill, pp. 183–190.
- Crystall B., Booth P. J., Klug D. R., Barber J. and Porter G. (1989), Resolution of a long-lived fluorescence component from D1/D2/cytochrome *b*559 reaction centres. *FEBS Lett.* **249**, 75–78.
- Danielius R. V., Satoh K., van Kan P. J. M., Plijter J. J., Nuijs A. M. and van Gorkom H. J. (1987), The primary reaction of photosystem II in the D1/D2/cytochrome *b*559 complex. *FEBS Lett.* **213**, 241–244.
- Eckert H.-J., Wiese N., Bernarding J., Eichler H.-J. and Renger G. (1988), Analysis of the electron transfer from Pheo⁻ to Q_A in PS II membrane fragments from spinach by time-resolved 325 nm absorption changes in the picosecond domain. *FEBS Lett.* **240**, 153–158.
- Khan E. M. (1993), An insight into the electronic and magnetic interactions in complexes of redox active ligands. Ph.D. Thesis, University of Poona, Pune (India).
- Klimov V. V., Allakhverdiev S. I. and Pashchenko V. Z. (1978), Measurement of activation energy and lifetime of fluorescence of photosystem II chlorophyll. *Dokl. Akad. Nauk. S.S.S.R.* (engl. transl.) **242**, 166–169.
- Klimov V. V., Allakhverdiev S. I., Demeter S. and Krasnovsky A. A. (1979), Pheophytin reduction in photosystem II of chloroplasts in relation to oxidation-reduction potential of medium. *Dokl. Akad. Nauk. S.S.S.R.* (engl. transl.) **249**, 190–192.
- Klimov V. V., Allakhverdiev S. I., Shuvalov V. A. and Krasnovsky A. A. (1982), Effect of extraction and re-addition of manganese on light reactions of photosystem II preparations. *FEBS Lett.* **148**, 307–312.
- Klimov V. V., Allakhverdiev S. I. and Ladygin V. G. (1986), Photoreduction of pheophytin in photosystem II of the whole cells of green algae and cyanobacteria. *Photosynth. Res.* **10**, 355–361.
- Marcus R. A. and Sutin N. (1985), Electron transfer in chemistry and biology. *Biochim. Biophys. Acta* **811**, 265–322.
- Michel H. and Deisenhofer J. (1988), Relevance of the photosynthetic reaction center from purple bacteria to the structure of photosystem II. *Biochemistry* **27**, 1–7.
- Moser C. C., Keske J. M., Warncke K., Farid R. S. and Dutton P. L. (1992), Nature of biological electron transfer. *Nature* **355**, 796–802.
- Nanba O. and Satoh K. (1987), Isolation of a photosystem II reaction center consisting of D1 and D2 polypeptides and cytochrome *b*559. *Proc. Natl. Acad. Sci. U.S.A.* **84**, 109–112.
- Nuijs A. M., van Gorkom H. J., Plijter J. J. and Duysens L. N. M. (1986), Primary-charge separation and excitation of chlorophyll *a* in photosystem II particles from spinach as studied by picosecond absorbance difference spectroscopy. *Biochim. Biophys. Acta* **848**, 167–175.
- Nixon R. J., Trost J. T. and Diner B. A. (1992), Role of the carboxy terminus of polypeptide D1 in the assembly of a functional water-oxidizing manganese cluster in photosystem II of the cyanobacterium *Synechocystis* sp. PCC 6803: assembly requires a free carboxyl group at C-terminal position 344. *Biochemistry* **31**, 10859–10871.
- Renger G. and Kaye A. (1987), On the mechanism of fluorescence quenching by photoaccumulating of the pheophytin anion radical in photosystem II. *Biochim. Biophys. Acta* **894**, 261–269.
- Renger G. (1992), Energy transfer and trapping in photosystem II, in: *Photosynthesis, The Photosystems: Structure, Function and Molecular Biology* (J. Barber, ed.). Elsevier, Amsterdam, pp. 45–99.
- Shuvalov V. A., Klimov V. V., Dolan E., Parson W. W. and Ke B. (1980), Nanosecond fluorescence and absorbance changes in photosystem II at low redox potential. *FEBS Lett.* **118**, 279–282.
- Shuvalov V. A., Heber U. and Schreiber U. (1989), Low temperature photochemistry and spectral properties of a photosystem II reaction center complex containing the proteins D1 and D2 and two heme of Cyt *b*559. *FEBS Lett.* **258**, 27–31.
- Takahashi Y., Hansson Ö., Mathis P. and Satoh K. (1987), Primary radical pair in the photosystem II reaction centre. *Biochim. Biophys. Acta* **893**, 49–59.
- Takahashi Y., Satoh K. and Itoh S. (1989), Silicomolybdate substitutes for the function of a primary electron acceptor and stabilizes charge separation in the photosystem II reaction center complex. *FEBS Lett.* **255**, 133–138.
- Trebst A. (1986), The topology of the plastoquinone and herbicide-binding peptides of photosystem II in the thylakoid membrane. *Z. Naturforsch.* **41c**, 240–245.
- Wiegardt K., Bossek U., Ventur D. and Weiss J. (1985), Assembly and structural characterization of binuclear μ -oxo-di- μ -acetato-bridged complexes of manganese(III) analogues of the diiron(III) centre in hemerythrin. *J. Chem. Soc. Chem. Commun.* 347–349.